

## Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

**Authors :** António Inês, Ana Guimarães, José Maria, Vânia Laranjo, Armando Venâncio, Luís Abrunhosa

**Abstract :** Lactic acid bacteria (LAB) play a key role in the biopreservation of a wide range of fermented food products, such as yogurt, cheese, fermented milks, meat, fish, vegetables (sauerkraut, olives and pickles), certain beer brands, wines and silage, allowing their safe consumption, which gave to these bacteria a GRAS (Generally Recognised as Safe) status. Besides that, the use of LAB in food and feed is a promising strategy to reduce the exposure to dietary mycotoxins, improving their shelf life and reducing health risks, given the unique mycotoxin decontaminating characteristic of some LAB. Mycotoxins present carcinogenic, mutagenic, teratogenic, neurotoxic and immunosuppressive effects over animals and Humans, being the most important ochratoxin A (OTA), aflatoxins (AFB1), trichothecenes, zearalenone (ZEA), fumonisin (FUM) and patulin. In a previous work of our group it was observed OTA biodegradation by some strains of *Pediococcus parvulus* isolated from Douro wines. So, the aim of this study was to enlarge the screening of the biodegradation over more mycotoxins besides OTA, including AFB1, and ZEA. This ability was checked in a collection of LAB isolated from vegetable (wine, olives, fruits and silage) and animal (milk and dairy products, sausages) sources. All LAB strains were characterized phenotypically (Gram, catalase) and genotypically. Molecular characterisation of all LAB strains was performed using genomic fingerprinting by MSP-PCR with (GTG)<sub>5</sub> and *csM13* primers. The identification of the isolates was confirmed by 16S rDNA sequencing. To study the ability of LAB strains to degrade OTA, AFB1 and ZEA, a MRS broth medium was supplemented with 2.0 µg/mL of each mycotoxin. For each strain, 2 mL of MRS supplemented with the mycotoxins was inoculated in triplicate with 10<sup>9</sup> CFU/mL. The culture media and bacterial cells were extracted by the addition of an equal volume of acetonitrile/methanol/acetic acid (78:20:2 v/v/v) to the culture tubes. A 2 mL sample was then collected and filtered into a clean 2 mL vial using PP filters with 0.45 µm pores. The samples were preserved at 4 °C until HPLC analysis. Among LAB tested, 10 strains isolated from milk were able to eliminate AFB1, belonging to *Lactobacillus casei* (7), *Lb. paracasei* (1), *Lb. plantarum* (1) and 1 to *Leuconostoc mesenteroides*. Two strains of *Enterococcus faecium* and one of *Ec. faecalis* from sausage eliminated ZEA. Concerning to strains of vegetal origin, one *Lb. plantarum* isolated from elderberry fruit, one *Lb. buchnerii* and one *Lb. parafarraginis* both isolated from silage eliminated ZEA. Other 2 strains of *Lb. plantarum* from silage were able to degrade both ZEA and OTA, and 1 *Lb. buchnerii* showed activity over AFB1. These enzymatic activities were also verified genotypically through specific gene PCR and posteriorly confirmed by sequencing analysis. In conclusion, due the ability of some strains of LAB isolated from different sources to eliminate OTA, AFB1 and ZEA one can recognize their potential biotechnological application to reduce the health hazards associated with these mycotoxins. They may be suitable as silage inoculants or as feed additives or even in food industry.

**Keywords :** bio-detoxification, lactic acid bacteria, mycotoxins, food and feed

**Conference Title :** ICNFS 2015 : 18th International Conference on Nutrition and Food Sciences

**Conference Location :** Zurich, Switzerland

**Conference Dates :** July 29-30, 2015

# BIODETOXIFICATION OF MYCOTOXINS BY LACTIC ACID BACTERIA FROM DIFFERENT FOOD MATRICES

António Inês<sup>1,2\*</sup>, Ana Guimarães<sup>3</sup>, Maria, J.C.<sup>1</sup>, Laranjo, V.<sup>1</sup>, A. Venâncio<sup>3</sup>, Luís Abrunhosa<sup>3</sup>

<sup>1</sup>Chemical Research Centre - Vila Real (CQ-VR), University of Trás-os-Montes and Alto Douro, Department of Biology and Environment, School of Life Sciences and Environment, Edifício de Enologia, Apartado 1013, 5001-801 Vila Real, Portugal. <sup>2</sup>Chemical Research Centre - Vila Real (CQ-VR), Chemistry Department, University of Trás-os-Montes and Alto Douro, School of Life Sciences and Environment, Vila Real, Portugal. <sup>3</sup>CEB-Centre of Biological Engineering, University of Minho, Campus de Gualtar, 4710-057, Braga, Portugal. \*aines@utad.pt

Lactic acid bacteria LAB play a key role in the biopreservation of a wide range of fermented food products, such as yogurt, cheese, fermented milks, meat, fish, vegetables (sauerkraut, olives and pickles), certain beer brands, wines and silage [1], allowing their safe consumption, which gave to these bacteria a GRAS (Generally Recognised as Safe) status. Besides that, the use of LAB in food and feed is a promising strategy to reduce the exposure to dietary mycotoxins, improving their shelf life and reducing health risks, given the unique mycotoxin decontaminating characteristic of some LAB. Mycotoxins present carcinogenic, mutagenic, teratogenic, neurotoxic and immunosuppressive effects over animals and Humans [2], being the most important ochratoxin A (OTA), aflatoxins (AFB1), trichothecenes, zearalenone (ZEA), fumonisin and patulin. In a previous work of our group it was observed OTA biodegradation by some strains of *Pediococcus parvulus* isolated from Douro wines [3].

So, the aim of this study was to enlarge the screening of the biodegradation over more mycotoxins besides OTA, including AFB1, and ZEA. This ability was checked in a collection of LAB isolated from vegetable (wine, olives, fruits and silage) and animal (milk and dairy products, sausages) sources. All LAB strains were characterized phenotypically (Gram, catalase) and genotypically. Molecular characterisation of all LAB strains was performed using genomic fingerprinting by MSP-PCR with (GTG)<sub>5</sub> and csM13 primers. The identification of the isolates was confirmed by 16S rDNA sequencing.

To study the ability of LAB strains to degrade OTA, AFB1 and ZEA, a MRS broth medium that was supplemented with 2.0 µg/mL of each mycotoxin was prepared. For each strain, 2 mL of MRS supplemented with the mycotoxins was inoculated in triplicate with 10<sup>9</sup> CFU/mL. The culture media and bacterial cells were extracted by the addition of an equal volume of acetonitrile/methanol/acetic acid (78:20:2 v/v/v) to the culture tubes. A 2 mL sample was then collected and filtered into a clean

2 mL vial using PP filters with 0.45 µm pores. The samples were preserved at 4 °C until HPLC analysis performed according to Abrunhosa e Venâncio [4]. Among LAB tested, 10 strains isolated from milk were able to eliminated AFB1, belonging to *Lactobacillus casei* (7), *Lb. paracasei* (1), *Lb. plantarum* (1) and 1 to *Leuconostoc mesenteroides*. Two strains of *Enterococcus faecium* and one of *Ec. faecalis* from sausage eliminated ZEA. Concerning to strains of vegetal origin, one *Lb. plantarum* isolated from elderberry fruit, one *Lb. buchnerii* and one *Lb. parafarraginis* both isolated from silage eliminated ZEA. Other 2 strains of *Lb. plantarum* from silage were able to degrade both ZEA and OTA, and 1 *Lb. buchnerii* showed activity over AFB1.


In conclusion, due the ability of some strains of LAB isolated from different sources to eliminate OTA, AFLA and ZEA one can recognize their potential biotechnological application to reduce the health hazards associated with these mycotoxins. They may be suitable as silage inoculants or as feed additives or even in food industry.

## References

- [1] M.E. Stiles, 1996. Biopreservation by lactic acid bacteria. *Antonie Van Leeuwenhoek*, 70 (1996), pp. 331–345
- [2] J.W. Bennett, M. Klich 2003. Mycotoxins. *Clin. Microbiol. Rev.*, 16 (2003), pp. 497–516.
- [3] Abrunhosa L, Inês A, Rodrigues AI, Guimarães A, Pereira VL, Parpot P, Mendes-Faia A, Venâncio A. 2014. *International Journal of Food Microbiology*, 188, 1: 45-52.
- [4] L. Abrunhosa, A. Venâncio, 2007. Isolation and purification of an enzyme hydrolyzing ochratoxin A from *Aspergillus niger*. *Biotechnology Letters*, 2007, 29, 1909-1914.

## Acknowledgements

This work was funded by FEDER funds through the COMPETE and by national funds through FCT, Ref. FCOMP-01-0124-FEDER-028029 and PTDC/AGR-TEC/3900/2012, respectively. Luís Abrunhosa received support through grant Incentivo/EQB/LA0023/2014 from ON.2 – O Novo Norte.




Zurich, Switzerland  
July 29 - 30, 2015

ICFSNPH 2015

17th International Conference on Food Safety, Nutrition and Public Health

# Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

António Inês, Ana Guimarães, José Maria, Vânia Laranjo, Armando Venâncio, Luís Abrunhos



Zurich, Switzerland  
July 29 - 30, 2015

ICFSNPH 2015

Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices


## INTRODUCTION

Lactic acid bacteria (LAB) play a key role in the biopreservation of a wide range of fermented food products, such as yogurt, cheese, fermented milks, meat, fish, vegetables (sauerkraut, olives and pickles), certain beer brands, wines and silage



allowing their safe consumption

which gave to these bacteria a GRAS (Generally Recognised as Safe) status




Zurich, Switzerland  
July 29 - 30, 2015

ICFSNPH 2015

Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## INTRODUCTION

Besides that, the use of LAB in food and feed is a promising strategy to reduce the exposure to dietary mycotoxins, improving their shelf life and reducing health risks, given the unique mycotoxin decontaminating characteristic of some LAB



Zurich, Switzerland  
July 29 - 30, 2015


ICFSNPH 2015

Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## INTRODUCTION

Mycotoxins present carcinogenic, mutagenic, teratogenic, neurotoxic and immunosuppressive effects over animals and Humans, being the most important ochratoxin A (OTA), aflatoxins (AFB1), trichothecenes, zearalenone (ZEA), fumonisin (FUM) and patulin

In a previous work of our group it was observed OTA biodegradation by some strains of *Pediococcus parvulus* isolated from Douro wines



Zurich, Switzerland  
July 29 - 30, 2015

ICFSNPH 2015

Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## OBJECTIVE

So, the aim of this study was to enlarge the screening of the biotransformation over more mycotoxins besides OTA, including AFB1, and ZEA. This ability was checked in a collection of LAB isolated from vegetable (wine, olives, fruits and silage) and animal (milk and dairy products, sausages) sources




Zurich, Switzerland  
July 29 - 30, 2015

ICFSNPH 2015

Bio-Detoxification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## MATERIAL AND METHODS

Global strategy for species identification and characterization of the LAB collection






ICFSNPH 2015

Bio-Defatification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## RESULTS AND CONCLUSION

Identification at species level and strain characterization of the lactic acid bacteria collection by phenotypic and genotypic approach





ICFSNPH 2015

Bio-Defatification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## RESULTS AND CONCLUSION

Source	Species	Mycotoxins
Milk	<i>Lactobacillus casei</i> (7)	AFB1
	<i>Lb. paracasei</i> (1)	
	<i>Lb. plantarum</i> (1)	
	<i>Leuconostoc mesenteroides</i> (1)	
Sausage	<i>Enterococcus faecium</i> (1)	ZEA
	<i>En. faecalis</i> (1)	
Elderberry fruit	<i>Lb. plantarum</i> (1)	ZEA
	<i>Lb. buchnerii</i> (1)	
Silage	<i>Lb. parafarraginis</i> (1)	ZEA and OTA
	<i>Lb. plantarum</i> (2)	
	<i>Lb. buchnerii</i> (1)	

OTA degradation:  
Carboxypeptidase




ICFSNPH 2015

Bio-Defatification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## RESULTS AND CONCLUSION

In conclusion, due the ability of some strains of LAB isolated from different sources to eliminate OTA, AFB1 and ZEA one can recognize their potential biotechnological application to reduce the health hazards associated with these mycotoxins

They may be suitable as silage inoculants or as feed additives or even in food industry



ICFSNPH 2015

Bio-Defatification of Mycotoxins by Lactic Acid Bacteria from Different Food Matrices

## Acknowledgments

This work was funded by FEDER funds through the COMPETE and by national funds through FCT, Ref. FCOMP-01-0124-FEDER-028029 and PTDC/AGR-TEC/3900/2012, respectively.

**FCT**  
Fundação para a Ciência e a Tecnologia  
Ministério da Educação e Ciência

Governo da República Portuguesa

**COMPETE**  
PROGRAMA OPERACIONAL FACTORES DE COMPETITIVIDADE

**QR**  
QUADRO DE REFERÊNCIA ESTRATÉGICO NACIONAL  
2007-2013

**WASET**  
WORLD ACADEMY OF SCIENCE, TECHNOLOGY AND ENVIRONMENT

**Universidade do Minho**

**CENTRE OF BIOLOGICAL ENGIN**

**IBB**  
INSTITUTE FOR BIOTECHNOLOGY AND BIOMANUFACTURING

**CQ**

**utad**  
UNIVERSIDADE DE TRÁS-OS-MONTES E ALTO DOURO